

Search for a Heavy SM Higgs Boson in the $H \rightarrow ZZ \rightarrow llqq$ Channel at ATLAS

Carl Gwilliam



UNIVERSITY OF
LIVERPOOL


(on behalf of the ATLAS Collaboration)




29th July 2011
Higgs Hunting 2011, Orsay

Introduction


- For a high mass Higgs ($m_H > 200$ GeV) ZZ and WW decay modes dominate
- “Golden” $H \rightarrow ZZ \rightarrow 4l$ is very clean but suffers from low branching fraction
- $H \rightarrow ZZ \rightarrow llqq$ has larger background but benefits from significantly higher BF
- Present the sensitivity of the ATLAS detector in this channel for 1.04 fb^{-1} at $\sqrt{s} = 7$ TeV in the range $200 \leq m_H \leq 600$ GeV



gluon fusion
(dominant)





vector boson
fusion (10-20%)



Signal modelled by NLO POWHEG MC generator interfaced to PYTHIA



Selection

- Leptonic Z candidate
 - 2 good isolated leptons (e/μ) with $p_T > 20$ GeV and $|\eta| < 2.5$
 - $76 < m_{ll} < 106$ GeV
- $E_T^{\text{miss}} < 50$ GeV
- Hadronic Z candidate
 - ≥ 2 jets with $p_T > 25$ GeV & $|\eta| < 2.5$
 - $70 < m_{jj} < 105$ GeV
- At high m_H the Z bosons from the H decay are boosted \rightarrow Additional cuts:
 - $P_T^{\text{jet}} > 45$ GeV
 - $\Delta\phi_{ll} < 90^\circ$ and $\Delta\phi_{jj} < 90^\circ$





Selection (2)

- Split into two samples based on identification of jets from b decays
 - Based on a combination of secondary vertex reconstruction and impact parameter significance of tracks within jet to primary vertex $\rightarrow \epsilon_b \approx 70\%$
- “Tagged”
 - Events with exactly 2 b -jets
 - Form invariant mass from the 2 tagged jets
- “Untagged”
 - Events with < 2 b -jets
 - Form invariant mass from all combinations of 3 leading jets





Z+jets Background

- Z+jets is the dominant background and is constrained using the m_{qq} sidebands
 - $40 < m_{qq} < 70$ GeV and $105 < m_{qq} < 150$ GeV
- Shape well described by ALPGEN but normalisation $\approx 10\%$ high
 - Consistent between tagged and untagged samples
- Use control region to determine scale factors to normalise MC



Top Background

- Top is an important background, particularly in the tagged sample
- Cross-checked using the sidebands of the m_{ll} distribution
 - $60 < m_{ll} < 76$ GeV and $106 < m_{ll} < 150$ GeV
 - For untagged sample also reverse E_T^{miss} cut
- Good description by MC@NLO Monte Carlo within errors





- Other backgrounds: QCD multijet production, also determined from data, and ZZ/WZ production, which are taken from MC@NLO



Results

- No significant excess over SM background observed

untagged



tagged



Systematic Uncertainties

● Signal

- Cross section uncertainty: 15 – 20% for $gg \rightarrow H$ & 3 – 9% for $qq \rightarrow qqH$
- Acceptance uncertainty by comparing POWHEG and PYTHIA

● Background normalisation

- Z+jets uncertainty from comparing low and high m_{qq} sidebands \rightarrow 1.4%/8.1% for low/high m_H untagged sample and 18% for tagged sample
- 100% uncertainty for QCD multijet and 50% for W+jets
- Theoretical uncertainty for top (9%) and ZZ/WZ (11%)

● Background shape

- Z+jets: comparison between ALPGEN and PYTHIA
- ZZ: comparison between MC@NLO and PYTHIA


● Luminosity: 3.7% (where normalisation not determined from data)

● Detector-related uncertainties on efficiency, E or p scale & resolution

- Tagged: dominated by uncertainty on b -tagging efficiency (15-25%)
- Untagged: Largest contribution is jet E scale (up to $\approx 5\%$) but jet E resolution, E_T^{miss} and b -tagging uncertainties also important

Exclusion Limits

- Combined exclusion limit on σ/σ_{SM} from tagged and untagged samples at 95% CL using *CLs* method
 - Use modified frequentist formalism with profile likelihood test statistic
 - The likelihood compares the full m_{lljj} distribution bin-by-bin to expected background or sum of expected signal and background
- Limit approaching σ_{SM} with $\int Ldt = 1.04 \text{ fb}^{-1}$
- Exclude $1.7 \times \sigma_{SM}$ at $m_H = 360 \text{ GeV}$
 - Corresponding expected limit is $2.7 \times \sigma_{SM}$
- Combined high mass $H \rightarrow ZZ/WW$ channels sensitive to σ_{SM}








† Limits based on on-shell Higgs with ad-hoc Breit-Wigner line shape. Potentially important effects related to off-shell Higgs boson production and interference effects are not included and may affect the limits for $m_H > 400 \text{ GeV}$

Conclusion



- Have presented the sensitivity of the ATLAS detector in the $H \rightarrow ZZ \rightarrow 2l2q$ channel with 1.04 fb^{-1} at $\sqrt{s} = 7 \text{ TeV}$
- No evidence for an excess above SM expectation observed
- Sensitivity ranges between 1.7 and 13 times SM cross section in the range $200 \leq M_H \leq 600 \text{ GeV}$, with maximal sensitivity at 360 GeV
- Channel contributes significantly to the combined exclusion limit in the high mass range
- LHC and ATLAS are performing well so expect improved sensitivity very soon

E_T^{miss} and b -tagging





$\Delta\phi_{ll}$ and $\Delta\phi_{jj}$



Results Table

ATLAS Preliminary

The expected number of signal and background candidates for the Higgs boson search for 1.04 fb^{-1} in the $H \rightarrow ZZ \rightarrow \ell\ell qq$ channel, along with the observed numbers of candidates in data. Numbers with uncertainties are \pm (stat.) \pm (syst.), respectively, and the statistical component assumes Gaussian uncertainties.

	Untagged			Tagged								
	Low- m_H		High- m_H	Low- m_H		High- m_H						
Z+jets	10352.	$\pm 61.$	$\pm 155.$	423.	$\pm 12.$	$\pm 30.$	72.	$\pm 1.$	$\pm 15.$	4.9	± 0.2	± 1.0
W+jets	10.	$\pm 2.$	5.	0.2	± 0.2	± 0.1	0.0	± 0.0	± 0.0	0.0	± 0.0	± 0.0
Top	40.	$\pm 1.$	6.	3.0	± 0.3	± 0.6	13.	$\pm 1.$	$\pm 3.$	1.1	± 0.2	± 0.3
Multijet	64.	$\pm 3.$	64.	2.0	± 0.5	± 2.0	0.3	± 0.2	± 0.3	0.0	± 0.0	± 0.0
ZZ	107.	$\pm 4.$	15.	8.5	± 1.1	± 1.8	6.9	± 1.0	± 2.0	0.79	± 0.23	± 0.30
WZ	143.	$\pm 3.$	29.	17.	$\pm 1.$	$\pm 3.$	0.5	± 0.2	± 0.3	0.03	± 0.02	± 0.01
Total background	10718.	$\pm 62.$	$\pm 173.$	453.	$\pm 13.$	$\pm 31.$	92.	$\pm 1.$	$\pm 15.$	6.9	± 0.4	± 1.2
Data	10495			419			91			6		
Signal												
$m_H = 200 \text{ GeV}$	33.	$\pm 1.$	6.				2.2	± 0.2	± 0.6			
$m_H = 300 \text{ GeV}$				7.0	± 0.3	± 1.5				0.58	± 0.08	± 0.19
$m_H = 400 \text{ GeV}$				9.8	± 0.3	± 1.8				1.1	± 0.1	± 0.3
$m_H = 500 \text{ GeV}$				5.5	± 0.1	± 1.0				0.63	± 0.04	± 0.19
$m_H = 600 \text{ GeV}$				2.5	± 0.1	± 0.5				0.28	± 0.02	± 0.08