


Probing fundamental physics with the Cosmic Microwave Background & Large Scale Structures



Anna Mangilli


Institut d'Astrophysique de Paris, IAP




LAL 25 February 2014



The Universe's history



The observable Universe




Concordance Λ CDM cosmological model

The questions list for Cosmology

Λ CDM

reference model =



what is the universe made of?


what is the nature of dark energy?

what is the nature of dark matter?


what is Inflation?

Where to search for answers

The Cosmic Microwave Background (CMB)




Large scale structures (LSS)





Primordial quantum fluctuations!

Planck



CMB Telescopes



LSS galaxy surveys

New observational data from CMB and galaxy surveys allow for precision tests of Λ CDM model and beyond!


Outline

- Probing late time evolution and primordial physics with the Cosmic Microwave Background (CMB) non-Gaussianity (NG)
 - Why CMB non-Gaussianity
 - Primordial and “late time” non-Gaussian signals
 - Planck Data analysis and future prospect
- CMB, Large scale structure and initial conditions
 - Constraining the nature of primordial perturbations beyond the Λ CDM model
 - Implications for CMB and LSS
 - Euclid+Planck forecasts


Why looking for non-Gaussianity (NG) in the CMB?

STANDARD INFLATIONARY MODEL predicts **GAUSSIAN CMB anisotropies**

If non-Gaussian signal in the CMB



PRIMORDIAL NG




DIFFERENT SCENARIO
FOR GENERATION
PRIMORDIAL
PERTURBATIONS?

The CMB non-Gaussianity is a very **high precision test**
of standard inflation!

Complementary to the search for CMB B-modes and power spectrum analysis

“LATE-TIME” NG



Primordial non-Gaussianity: an example



Non-linear gravitational potential perturbations

$$\Phi(\mathbf{x}) = \Phi_L(\mathbf{x}) + f_{NL}(\Phi_L^2(\mathbf{x}) - \langle \Phi_L^2(\mathbf{x}) \rangle)$$

Salopek & Bond 1990, Gangui et al. 1994
Verde et al. 2000, Komatsu & Spergel 2001



AMPLITUDE of the quadratic non-linear correction

Small for standard slow roll inflation, large for models e.g. multi field inflation

Different NG phenomena leave different imprints in the CMB sky which can be used to constrain the physical mechanism behind them.

The “late-time” CMB non-Gaussianity

Uncorrelated CMB photons



ISW=Integrated Sachs Wolfe

$$\dot{\Phi} \neq 0$$

Non-gaussian signal in the CMB
due to the cross correlation two effects:

ISW – CMB photon red/blue shifted: dark energy stretches the gravitational potential wells


LENSING – CMB photon deflected by the forming structures

The CMB lensing-ISW non-Gaussianity

Direct probe of the action of Dark Energy on the evolution of structures

The lensing-ISW biases the primordial NG

Contamination of primordial local non-Gaussianity due to the late time signal



Mangilli&Verde 2009



**BIAS to the primordial signal: Δf_{NL} of order 10, bigger than
Planck 1- σ error on primordial f_{NL}**


How look for non-Gaussianity in the CMB

2-points correlation function

$$\left\langle \frac{\Delta T}{T}(\mathbf{x}_1) \frac{\Delta T}{T}(\mathbf{x}_2) \right\rangle$$

$$\frac{\Delta T(\mathbf{n})}{T_0} = \sum_{\ell=0}^{\infty} \sum_{m=-\ell}^{\ell} a_{\ell m}^T Y_{\ell m}(\mathbf{n})$$

Angular power spectrum





Non-Gaussianity? More information!

Look at the higher order statistics
beyond the power spectrum

3-points correlation function

$$\left\langle \frac{\Delta T}{T}(\mathbf{x}_1) \frac{\Delta T}{T}(\mathbf{x}_2) \frac{\Delta T}{T}(\mathbf{x}_3) \right\rangle$$

Angular bispectrum




Look at
triangles in the
CMB!

Different mechanisms, different amplitudes and shapes!

$$B(\ell_1, \ell_2, \ell_3) = f_{\text{NL}} F(\ell_1, \ell_2, \ell_3)$$

SHAPE



SQUEEZED shape


Primordial (local type):

- ▶ Multi-fields inflation
- ▶ Ekpyrotic/cyclic models
- ▶ Curvaton isocurvature model


Non-primordial:

- ▶ Lensing-ISW correlation

ISW-I.




EQUILATERAL shape



- ▶ non-canonical kinetic term
- ▶ Dirac-Born-Infeld (DBI) inflation
- ▶ Ghost inflation

ORTHOGONAL shape

- ▶ non-canonical kinetic term
- ▶ higher derivatives interactions





Planck data analysis and results on CMB non-Gaussianity

On behalf of the Planck collaboration



The Planck experiment

Planck is an ESA mission which observed the sky in 9 frequency bands from 30 to 857 GHz with an unprecedented sensitivity




Component separation to remove foregrounds.


Cleaned Planck CMB map

The Planck bispectrum


SMICA



NILC




SEVEM




Robust to foreground cleaning

Constraints on fNL from Planck data



Local squeezed



Equilateral



Orthogonal


KSW

Binned

Modal

SMICA

	KSW	Binned	Modal
Local	9.8 ± 5.8	9.2 ± 5.9	8.3 ± 5.9
Equilateral	-37 ± 75	-20 ± 73	-20 ± 77
Orthogonal	-46 ± 39	-39 ± 41	-36 ± 41



Lensing-ISW bias

KSW	7.7 ± 1.5
Binned	7.7 ± 1.6
Modal	10 ± 3

Planck Collaboration 2013


Mangilli&Verde 2009




ISW-lensing subtracted

KSW	Binned	Modal
2.7 ± 5.8	2.2 ± 5.9	1.6 ± 6.0
-42 ± 75	-25 ± 73	-20 ± 77
-25 ± 39	-17 ± 41	-14 ± 42

No evidence of primordial NG in Planck Data




Planck results on the lensing-ISW



Bispectrum template

Planck data (SMICA)



**Specific estimators calibrated
for the L-ISW signal**

Mangilli et. al 2013

Planck Collaboration 2013

Expected amplitude LCDM

$$f_{\text{NL}}^{\text{ISW-L}} = 1$$

Amplitude of the lensing-ISW bispectrum $f_{\text{NL}}^{\text{ISW-L}}$

Estimator	NILC	SEVEM	SMICA
$T\phi$	0.75 ± 0.28	0.62 ± 0.29	0.70 ± 0.28
KSW	0.85 ± 0.32	0.68 ± 0.32	0.81 ± 0.31
binned	1.03 ± 0.37	0.83 ± 0.39	0.91 ± 0.37
modal	0.93 ± 0.37	0.60 ± 0.37	0.77 ± 0.37

Consistent results from all estimators and data maps with different component separation methods



**Planck finds evidence for the first time of the
Lensing-ISW signal at 2.7σ**



Planck high resolution!



Primordial non-Gaussianity: Local shape



f_{NL}^{local} WMAP9 = 37 ± 20

Bennet et al. 2012

f_{NL}^{local} Planck 2013 = 9.8 ± 5.8


The Planck collaboration 2013

Consistency with WMAP

Polarization forecasts

For primordial non-Gaussianity:

PLANCK T+Pol



For lensing-ISW non-Gaussianity:

	$\sigma_{f_{NL}}$	σ_{lens}	correlation	bias on f_{NL}	$\sigma_{f_{NL}}^{\text{marge}}$
T	4.31	0.19	0.24	9.5	4.44
T+E	2.14	0.12	0.022	2.6	2.14
Planck T	5.92	0.26	0.22	6.4	6.06
Planck T+E	5.19	0.22	0.13	4.3	5.23

Lewis et al. 2011

T+Pol $\sim 15\%$ improvement

CMB non-Gaussianity: TAKE AWAY message!

- **Non-Gaussianity in the CMB: powerful tool to constrain primordial physics and Dark Energy (late-time lensing-ISW bispectrum)**
- **Planck constrained for the first time CMB Non-Gaussianity with unprecedented precision!**



Planck favors the simplest models for inflation



Planck finds evidence for the first time of the Integrated-Sachs-Wolfe-lensing bispectrum. Signal compatible with the LCDM scenario

-

Future prospects:



Planck polarization data 2014 release!



Lensing-ISW bispectrum: new observable to be used to constrain dark energy properties.


Outline

- Probing late time evolution and primordial physics with the Cosmic Microwave Background (CMB) non-Gaussianity (NG)
 - Why CMB non-Gaussianity
 - Primordial and “late time” non-Gaussian signals
 - Planck Data analysis and future prospect
- CMB, Large scale structure and initial conditions
 - Constraining the nature of primordial perturbations beyond the Λ CDM model
 - Implications for CMB and LSS
 - Euclid+Planck forecasts


Standard model for structure formation

standard INFLATION: single scalar field, the inflaton, drives accelerated expansion **and** produce primordial perturbations

Cosmic inflation: Origin of the perturbations



What is the nature of the primordial fluctuations?



Cosmic inflation: Origin of the perturbations


Curvature adiabatic perturbation

[Standard Inflationary dynamics](#) implies that constant density perturbations are present initially. Perturbations in all components are spatially homogeneous.

Isocurvature entropy perturbation

No initial curvature perturbations. Fluctuation in number density between different components. The initial density fluctuations are created from stresses in the radiation–matter component. E.g. Cold Dark Matter and Neutrinos Isocurvature modes.


New observational data from CMB and galaxy surveys offers precision tests of the nature of the primordial perturbations




Photon and matter decoupling: CMB

Dark matter + baryon and photons tightly coupled

BARYON ACOUSTIC OSCILLATIONS (BAO) = sound waves



BAO froze at decoupling leaving an imprint in the CMB: the characteristic peak structure in the CMB power spectrum!




Pure isocurvature ruled out but ...

Current observations allow for mixed Adiabatic+Isocurvature initial conditions

f_{iso}
Isocurvature/adiabatic ratio
parameter
95% CL upper bound

General model:		f_{iso}
CDM isocurvature	<u>dark matter</u>	0.39
ND isocurvature	<u>neutrinos</u>	0.27
NV isocurvature		0.14
Special CDM isocurvature cases:		
Uncorrelated, $n_{II} = 1$, ("axion")		0.039
Fully correlated, $n_{II} = n_{RR}$, ("curvaton")		0.0025
Fully anti-correlated, $n_{II} = -n_{RR}$		0.0087

The Planck collaboration 2013



✓ Allowing for **isocurvature** modes introduces new **degeneracies in the parameters space** which can compromise accuracy of parameters constraints (**systematic shifts and bias**)



Extra isocurvature contribution can bias CMB parameter estimation and modify the constraint of the typical scale of the Baryon Acoustic Oscillation imprinted in the CMB

BAO scale r_s = “standard ruler”




The BAO scale as imprinted in the CMB evolves as the Universe expands and remains also imprinted in the large scale matter distribution and can be used to probe Dark Energy by galaxy surveys



EUCLID

Baryon Acoustic Oscillations (BAO) scale r_s = standard ruler

in the LSS




Future surveys i.e . Euclid will be able to measure BAO scale with very high accuracy

Extra isocurvature contribution can bias CMB parameter estimation and modify the constraint of the typical scale of the Baryon Acoustic Oscillation imprinted in the CMB


BAO scale r_s = “standard ruler”

The BAO scale as imprinted in the CMB evolves as the Universe expands and remains also imprinted in the large scale matter distribution


Energy by galaxy surveys




Baryon Acoustic Oscillations (BAO)
scale r_s = standard ruler



in the LSS



EUCLID






A wrong assumption on the nature of primordial perturbations leads to a systematic **bias** on the BAO scale measurements from large scale structure surveys **bigger than survey experimental errors i.e EUCLID**

Wrong interpretation of Dark energy properties from galaxy surveys data

Combining CMB and Large scale structures

Combining the information from Large Scale Structure (LSS) survey and CMB breaks parameter degeneracies and greatly improves constraint on the nature of the primordial perturbations and on **Dark Energy parameters**

isocurvature fraction parameter



EUCLID + Planck (T&Pol)



Dark energy equation of state parameters

CMB and galaxy surveys allow for precision tests of the standard Λ CDM and beyond!

- Planck constrained for the first time CMB Non-Gaussianity with unprecedented precision!

Planck favors the simplest models for inflation

Planck finds evidence for the first time of the Integrated-Sachs-Wolfe-lensing bispectrum: new observable!

- Combining CMB + Large Scale Structure very powerful!

Degeneracies are solved (no systematic shifts) and constraints on the nature of primordial perturbations improved even for extended model

More data!

Planck full mission and **POLARISATION** data on 2014

Current research projects



2014 release

Planck non-Gaussianity WG

- New constraints on primordial and late time NG adding polarization data

Planck HFI likelihood WG group

- Planck temperature and polarization likelihood analysis for cosmological parameter estimation within the Λ CDM scenario and beyond



THANK YOU FOR YOUR ATTENTION!



BIBLIOGRAPHY:

Mangilli, Wandelt, Elsner, Liguori A&A 2013

Carbone, Mangilli, Verde JCAP 2011

Mangilli, Verde, Beltran JCAP 2010

Mangilli, Verde PRD 2009

Mangilli, Bartolo, Matarrese, Riotto PRD 2008

----- Planck results 2013, A&A 2013:

The Planck Collaboration (incl. Mangilli): XXIV. Constraints on primordial non Gaussianity

The Planck Collaboration (incl. Mangilli): XIX. The integrated Sachs-Wolfe effect

The Planck Collaboration (incl. Mangilli): XVII. Gravitational lensing by large-scale structure

The Planck Collaboration (incl. Mangilli): XVIII. Isotropy and statistics