



ID de Contribution: 25

Type: Talk

## Study of key resonances in the $^{30}\text{P}(p,g)^{31}\text{S}$ reaction in classical novae

*mardi 29 mai 2018 10:45 (20 minutes)*

Classical novae outbursts are the third most energetic explosions in the Universe after gamma-ray bursts and supernovae. During this explosive burning, nucleosynthesis takes place and the newly synthesized material is ejected into the interstellar medium. In order to understand these objects, the study of presolar grains and  $\gamma$ -ray emitters are of specific interest since they can give direct insights into the nucleosynthesis processes and isotopic abundances.

The  $^{30}\text{P}(p,g)^{31}\text{S}$  reaction is one of the few remaining reactions with a rate uncertainty which has a strong impact on classical novae model predictions. Sensitivity studies have shown that it has the largest impact on the predicted elemental abundance ratios of Si/H, O/S, S/Al, O/P and P/Al, which can be used to constrain physical properties of classical novae. The  $^{30}\text{Si}/^{28}\text{Si}$  isotopic ratio, which is an important signature that helps to identify presolar meteoritic grains of a likely nova origin, depends also strongly on the  $^{30}\text{P}(p,\gamma)^{31}\text{S}$  reaction rate.

To reduce the nuclear uncertainties associated to this reaction we performed an experiment at ALTO facility of Orsay using the  $^{31}\text{P}(^3\text{He},t)^{31}\text{S}$  reaction to populate  $^{31}\text{S}$  excited states of astrophysical interest. The tritons were momentum analysed using the Enge Split-Pole magnetic spectrometer and the decaying protons were detected in coincidence in an array of DSSSDs (Double Sided Silicon Stripped Detectors). The comparison of the focal plane spectra obtained for single and coincidence events will allow the extraction of the proton branching ratios.

After a presentation of the astrophysical context of this work, the current situation of the  $^{30}\text{P}(p,g)^{31}\text{S}$  reaction rate will be discussed. Then the experiment set up and the analysis of the single events from the Split-Pole focal plane detector will be presented.

**Auteur principal:** Mme MEYER, Anne (IPN Orsay)

**Orateur:** Mme MEYER, Anne (IPN Orsay)

**Classification de Session:** Nuclear Physics - Experiments