

# Cold muonium beam for atomic physics and gravity experiments

*mardi 15 octobre 2019 19:40 (5 minutes)*

We are investigating methods to create a novel muonium (Mu) source, based on  $\mu^+ \rightarrow \text{Mu}$  conversion in superfluid helium (SFHe), which has the potential of providing high brightness Mu beams for next generation laser spectroscopy experiments. We are also investigating the feasibility of using such sources for measuring the gravitational interaction of Mu. The positive muon ( $\mu^+$ ) which is dominating the Mu mass is not only an elementary antiparticle, but a second-generation lepton too. This makes a gravity experiment highly motivated, and complementary to gravitational studies of antihydrogen and positronium.

State-of-the-art Mu sources (like silica aerogel, mesoporous SiO<sub>2</sub>) emit Mu atoms with a large (thermal) energy distribution, and wide ( $\sim \cos \theta$ ) angular distribution. Cooling of these porous samples below 100 K results in rapidly declining numbers of vacuum-emitted muonium due to decreased mobility, and atoms sticking to the pore walls.

Our proposed method relies on stopping  $\mu^+$  in a thin layer of SFHe, and forming Mu by capturing an electron from the ionization trail. A fraction of the Mu diffuses to the SFHe surface within their lifetime, where emission into vacuum occurs. The velocity of the emitted Mu ( $\sim 6 \text{ mm}/\mu\text{s}$ ) is given by their large chemical potential ( $E/k_B \sim 270 \text{ K}$ ) in SFHe, while their thermal distribution determines the transverse momentum.

In this poster, methods and challenges to create such SFHe Mu sources, the present status, and the feasibility of an antimatter gravity experiment will be discussed.

**Orateur:** SOTER, Anna (Paul Scherrer Institute)

**Classification de Session:** Poster session