

# A NLO+PS generator for $H \rightarrow VV$ production in gluon fusion including non-resonant and offshell effects



Silvia Ferrario Ravasio  
Oxford University





## Higgs Hunting 2021

Orsay-Paris, online, 20<sup>th</sup> September 2021

based on [Eur. Phys. J. C 81, 687 \(2021\)](#), [[2102.07783](#)], S. Alioli, S.F.R., J.M. Lindert and R. [Röntsch](#)

# Anatomy of $gg \rightarrow H \rightarrow VV$

- Gluon fusion is the dominant mechanism for Higgs production at the LHC
- $H \rightarrow VV$  sensitive to the Higgs - gauge bosons coupling
- Roughly 10% of  $gg \rightarrow H \rightarrow VV$  comes from  $m_W > m_V$
- Offshell Higgs cross section important to determine  $\Gamma_H \ll$  detector resolutions
- **QCD background**  $gg \rightarrow VV$  is dominant and cannot be distinguished from the **signal**
- The **full** contribution is given by the sum of background, signal as well as their **interference**



Sizeable and negative **interference** for large  $m_W$

# Anatomy of $pp \rightarrow VV \rightarrow 4l$

- $gg \rightarrow VV$  contributes to the NNLO QCD corrections to  $pp \rightarrow VV$ , and can be computed separately

ATLAS fiducial cuts for  $gg \rightarrow ZZ \rightarrow 4l$  @ 13 TeV,  
1902.05892

Grazzini, Kallweit, Wieseemann, Yook '21

Contribution	$\sigma$ [fb]
LO	$36.8^{+2.9}_{-2.6}$
NLO	$49.0^{+1.5}_{-1.4}$
NNLO (no gg)	$52.1^{+0.7}_{-0.7}$
gg @ LO	$4.3^{+1.1}_{-0.8}$
gg @ NLO	$7.8^{+1.3}_{-1.1}$

$O(\alpha_s^2) = 3.1 + 4.3$  pb, the gluon-fusion channel is enhanced by the large gluon luminosity

Large NLO corrections



A lot of recent activity!

- $pp \rightarrow WW \rightarrow 4l$  ([MiNNLO<sub>ps</sub>](#), [Lombardi, Wieseemann, Zanderighi '21](#), [[Re, Wieseemann, Zanderighi '18](#)]) and  $pp \rightarrow ZZ \rightarrow 4l$  ([GENEVA](#), [Alioli, Broggio, Gavardi, Kallweit, Lim, Nagar, Napoletano '21](#); [MiNNLO<sub>ps</sub>](#); [Buonocore, Koole, Lombardi, Rottoli, Wieseemann, Zanderighi '21](#)) are both known at **NNLOPS**.
- In this talk:  $gg \rightarrow VV \rightarrow 4l$  at **NLOPS** in [POWHEG BOX RES](#), with spin correlations, interferences and off-shell effects are included exactly, top-quark mass effects are included approximately in the QCD bkgd ([S. Alioli, S.F.R., J.M. Lindert and R. Röntsch '21](#))

# gg4l @ NLO in POWHEG BOX RES

- First NLOPS for  $gg \rightarrow VV \rightarrow 4l$  including off-shell effects, **interference** between the Higgs mediated signal and the **QCD background**. Exact one-loop matrix elements from **OpenLoops** and two loops from **Caola etal, 2016** (and **ggVVamp**).
- **VV = WW**: **top-mass effects** for the virtual **bkgd** ampl obtained using LO reweighting (@ fixed helicity, **MCFM**)

$$\mathcal{A}_{\text{bkgd}}^{(2), WW} = \mathcal{A}_{\text{bkgd}}^{(2), WW}(u, d, s, c) \frac{\mathcal{A}_{\text{bkgd}}^{(1), WW}(u, d, s, c, b, \mathbf{t})}{\mathcal{A}_{\text{bkgd}}^{(1), WW}(u, d, s, c)}$$





- **VV = ZZ**: **top-mass effects** for the virtual **background** amplitude obtained using large- $m_t$  expansion from **Caola etal, 2016**. (LO rwgt also available for large  $m_{4l}$  region, as well as interpolation between the two options)

Top-mass effects in **bkgd** ampl important for the **interference**: offshell Higgs decays preferentially to longitudinal Z's, which couple more with top quark loops than with massless loops

# gg4l @ NLOPS in POWHEG BOX RES: setup



- pp collisions @ **13 TeV**
- Central renormalization and factorization scale  $\mu = m_{4l}/2$
- **ZZ** channel:
  - 5 GeV <  $m_{ll}$  < 180 GeV;
  - 10 GeV <  $m_{4l}$  < 340 GeV;
- Les Houches level (LHE) predictions by POWHEG matched to the **PYTHIA8.2** general purpose Monte Carlo event generator (**default shower, PowhegHooks** class to veto emissions harder than the POWHEG one)

# gg4l @ NLOPS in POWHEG + PYTHIA: $m_{ZZ}$ & $m_{TWW}$



Invariant (transverse) mass of the VV system left unchanged by the parton shower. The relative size of the **signal** and of its **interference** with the QCD **background** increases in the tail.


# gg4l @ NLOPS in POWHEG + PYTHIA: $H_T$



WW

Large impact of multiple PS emissions for the tail of  $H_T = \sum_{l,\nu,j} p_\perp$ , particularly for the **signal**, which at f.o. is peaked at smaller values.



# gg4l @ NLOPS in POWHEG + PYTHIA: $p_{Tj}$



Resummation effects in LHE and NLOPS distributions sizeable for small values of the hardest jet  $p_T$ , in the tail agreement with fixed order.



# gg4l @ NLOPS in POWHEG + PYTHIA: $p_{TVV}$




Z


WW

At NLO and LHE,  $p_{TVV} = p_{Tj}$ . The PS enhances significantly the tail, where scale variations are large.

# gg4l @ NLOPS in POWHEG + PYTHIA: $p_{T,4l}$



+50% PS corrections in the tail of  $p_{T,4l}$  depend on the PS recoil scheme. By **default**, transverse momentum imbalance due to ISR always absorbed by the final state. In the **Catani-Seymour shower**, if the incoming emitter is in an initial-final dipole, the final state spectator takes the recoil. Differences are at **next-to-leading log** accuracy and as large as scale variations.




## Summary and conclusions

- $gg \rightarrow H \rightarrow VV$  important to probe the HVV coupling: 10% of the cross-section comes from the region  $m_{VV} \gg m_V$ , which can be used also for  $\Gamma_H$  determinations.
- We have implemented in POWHEG BOX RES the **first NLOPS generator for off-shell VV production**, with leptonic V decay, in **gluon fusion** including the Higgs mediated **signal**, his QCD **background** and their **interference**.
- One loop matrix elements are exact, some approximations are made for the **top-mass** dependence of the **bkgd** amplitude at two loops: we can replace them once the exact calculation becomes available.
- We performed a phenomenological study at 13 TeV: **PS effects** are sizable e.g. for  $pT_{4l}$  and  $H_T$ .
- Large dependence on the PS recoil scheme in the tail of the  $pT_{4l}$  distributions, whose accuracy is only LO+LL.
- The code will be released in a couple of weeks:

`svn co --username anonymous --password anonymous svn://powhegbox.mib.infn.it/trunk/User-Processes-RES/gg4l`


# BACKUP

# gg4l @ NLO in POWHEG BOX RES




- Singular qq-initiated** and **regular qq-initiated** contributions are included in the real corrections and contribute to the 20% of the **total**

Sizeable impact  
(but scale variations unchanged)



Regular,  
negligible



# gg4l @ NLOPS in POWHEG BOX RES: setup

- pp collisions @ **13 TeV**
- **NNPDF31\_nlo\_as\_0118** PDF set
- Central renormalization and factorization scale  $\mu = m_{4l}/2$
- $m_b = 0$  (**5 flavour** scheme)  $m_t = 173.2$  GeV
- **ZZ** channel:  $5 \text{ GeV} < m_{ll} < 180 \text{ GeV}$ ;  $m_{4l} < 340 \text{ GeV}$
- **WW** channel:  $m_{2l2\nu} > 1 \text{ GeV}$
- **Jets**: anti-kt,  $R=0.4$ ,  $p_T > 20 \text{ GeV}$
- POWHEG matching with **bornzerdamp** and **hdamp=100 GeV** to separate the real contribution into singular and not singular

$$d\sigma = \left[ B(\Phi_b) + V(\Phi_b) + \int d\Phi_{\text{rad}} R_s(\Phi_b, \Phi_{\text{rad}}) \right] \left[ \Delta^{\text{PWG}}(p_{\perp \text{min}}) + \frac{R_s(\Phi_b, \Phi_{\text{rad}})}{B(\Phi_b)} \Delta^{\text{PWG}}(p_{\perp}(\Phi_{\text{rad}})) \right] + R_{ns}(\Phi_{b+1})$$

- Les Houches level (LHE) predictions by POWHEG matched to the **PYTHIA8.2** general purpose Monte Carlo event generator (**default shower**, **PowhegHooks** class to veto emissions harder than the POWHEG one)